
**Open-File Report 96-343** 

# Base-Flow Data for the Little West Fork Basin, Fort Campbell, Tennessee and Kentucky, 1993 and 1994



Prepared by the U.S. Geological Survey

in cooperation with the United States Department of the Army, Fort Campbell, Directorate of Public Works, Environmental Division



maintaining the data needed, and c including suggestions for reducing	ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding an	o average 1 hour per response, inclu- ion of information. Send comments arters Services, Directorate for Infor ny other provision of law, no person	regarding this burden estimate or mation Operations and Reports	or any other aspect of the property of the pro	nis collection of information, Highway, Suite 1204, Arlington
1. REPORT DATE		2. REPORT TYPE		3. DATES COVE	RED
1996		N/A		-	
4. TITLE AND SUBTITLE				5a. CONTRACT	NUMBER
Base-Flow Data for and Kentucky, 199		rk Basin, Fort Camp	pbell, Tennessee	5b. GRANT NUN	ИBER
and Kentucky, 199	3 anu 1994			5c. PROGRAM E	LEMENT NUMBER
6. AUTHOR(S)				5d. PROJECT NU	JMBER
				5e. TASK NUME	BER
				5f. WORK UNIT	NUMBER
	ZATION NAME(S) AND AE  f the Interior 1849 (	DDRESS(ES) C <b>Street, NW Washi</b>	ngton, DC	8. PERFORMING REPORT NUMB	G ORGANIZATION ER
9. SPONSORING/MONITO	RING AGENCY NAME(S) A	AND ADDRESS(ES)		10. SPONSOR/M	ONITOR'S ACRONYM(S)
				11. SPONSOR/M NUMBER(S)	ONITOR'S REPORT
12. DISTRIBUTION/AVAIL Approved for publ	LABILITY STATEMENT ic release, distributi	on unlimited			
13. SUPPLEMENTARY NO	TES				
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFIC	ATION OF:		17. LIMITATION OF	18. NUMBER	19a. NAME OF
a. REPORT unclassified	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE unclassified	ABSTRACT SAR	OF PAGES  34	RESPONSIBLE PERSON

**Report Documentation Page** 

Form Approved OMB No. 0704-0188

**Cover illustration.** Graphic by J.E. Banton and K.A. Carney; photograph by G.E. Hileman.

## Base-Flow Data for the Little West Fork Basin, Fort Campbell, Tennessee and Kentucky, 1993 and 1994

By David E. Ladd

U.S. GEOLOGICAL SURVEY

Open-File Report 96-343

Prepared in cooperation with the

UNITED STATES DEPARTMENT OF THE ARMY, FORT CAMPBELL, DIRECTORATE OF PUBLIC WORKS, ENVIRONMENTAL DIVISION



Nashville, Tennessee 1996

# U.S. DEPARTMENT OF THE INTERIOR BRUCE BABBITT, Secretary

U.S. GEOLOGICAL SURVEY Gordon P. Eaton, Director

Any use of trade, product, or firm name in this report is for identification purposes only and does not constitute endorsement by the U.S. Geological Survey.

For additional information write to:

District Chief U.S. Geological Survey 810 Broadway, Suite 500 Nashville, Tennessee 37203 Copies of this report may be purchased from:

U.S. Geological Survey
Earth Science Information Center
Open-File Reports Section
Box 25286, MS 517
Denver Federal Center
Denver, Colorado 80225

## **CONTENTS**

Abstr	ract	1
Introd	duction	1
Base-	-flow data	2
	Discharge data	
	Temperature data	18
	Specific-conductance data	
Areas	s of surplus or deficient flow	18
Sumr	mary	28
Refer	rences cited	28
FIGL	JRES	
1.	Map showing location of the study area and locations of stream and spring measurement sites	3
2.	Hydrograph showing mean daily discharge on the Little West Fork at site number 40 in Tennessee	Q Q
3-5.		Ü
	discharge measurements:	
	3. Low base-flow, September 1 and 13, 1993	11
	4. High base-flow, March 17 and 18, 1994	14
	5. Low base-flow, October 18 and 19, 1994	17
6-8.	Maps of the Little West Fork basin showing areas of surplus or deficient flow during:	- '
	6. Low base flow, September 1 and 13, 1993	25
	7. High base flow, March 17 and 18, 1994	26
	8. Low base flow, October 18 and 19, 1994	27
TABL	LES	
1.	Site and station numbers, station names, and locations of stream and spring measurement sites	
	in the Little West Fork basin, Fort Campbell	4
2.	Low base-flow data for streams in the Little West Fork basin, September 1 and 13, 1993	
3.	Low base-flow data for springs in the Little West Fork basin, September 13, 1993	10
4.	High base-flow data for streams in the Little West Fork basin, March 17 and 18, 1994	
5.	High base-flow data for springs in the Little West Fork basin, March 17, 1994	
6.	Low base-flow data for streams in the Little West Fork basin, October 18 and 19, 1994	
	Low base-flow data for springs in the Little West Fork basin, October 18, 1994	
8.	Low base-flow data for drainage areas in the Little West Fork basin, September 1 and 13, 1993	19
9.	High base-flow data for drainage areas in the Little West Fork basin, March 17 and 18, 1994	21
	Low base-flow data for drainage areas in the Little West Fork basin, October 18 and 19, 1994	
	Flow per square mile for major tributary basins in the Little West Fork basin	
		- •

#### **CONVERSION FACTORS AND ABBREVIATIONS**

Multiply	Ву	To obtain
inch (in.)	25.4	millimeter
foot (ft) mile (mi)	0.3048 1.609	meter kilometer
square mile (mi <sup>2</sup> ) cubic foot per second (ft <sup>3</sup> /s)	2.59 0.02823	square kilometer cubic meter per second
cubic foot per second per square mile (ft <sup>3</sup> /s/mi <sup>2</sup> )	0.01090	cubic meter per second per square kilometer
microSiemen per centimeter (μS/cm)	1	micromho per centimeter

Temperature in degrees Fahrenheit (°F) can be converted to degrees Celsius (°C), and temperature in °C to °F, as follows:  $^{\circ}F = 1.8^{\circ}C + 32$   $^{\circ}C = 5/9(^{\circ}F - 32)$ 

# Base-Flow Data for the Little West Fork Basin, Fort Campbell, Tennessee and Kentucky, 1993 and 1994

By David E. Ladd

#### **Abstract**

Base-flow data were collected from selected sites within the Little West Fork basin during high and low base-flow conditions to support a study of the source and movement of ground water that supplies the Fort Campbell Military Reservation. Stream and spring discharge, water temperature, and specific-conductance data were collected during low base-flow conditions from 64 sites on September 1 and 13, 1993, and again on October 18 and 19, 1994. High baseflow data were collected from 63 sites on March 17 and 18, 1994. Discharge was greater during high base-flow conditions than during low base-flow conditions. Major tributaries on the south side of the study area consistently had lower flow than the tributaries on the north side.

Discharge data were used to categorize stream reaches and sub-basins. Stream reaches were categorized as gaining or losing, wet, dry, or unobserved for each base-flow measurement period. More gaining stream reaches occurred during the high base-flow period than either of the low base-flow periods. More dry stream reaches occurred during the two low base-flow periods than during the high base-flow period.

Sub-basin areas with surplus or deficient flow were also defined. Many areas of deficient flow occurred near the headwaters of the Little West Fork basin under all base-flow conditions. Fewer areas of deficient flow occurred near the mouth of the basin.

The flow per square mile for each major tributary basin in the study area was also calculated. The values of flow per square mile for the

Noah Spring Branch and Dry Fork Creek basins in the northern part of the study area were greater than those for the Piney Fork and Fletchers Fork basins in the southern part of the study area under all base-flow conditions.

#### INTRODUCTION

In recent years, environmental managers at the Fort Campbell Military Reservation have become increasingly concerned with the need for protection and management of the Reservation's ground-water resources. In 1993, the U.S. Geological Survey (USGS), in cooperation with the U.S. Army and Fort Campbell, began a study to better understand the occurrence and movement of ground water in the Fort Campbell area. As part of this study, a base-flow investigation of the drainage basin for the Little West Fork of the Red River was conducted. Base-flow data including stream and spring discharge, water temperature, and specific conductance were collected from selected sites within the basin during 1993 and 1994. Data were collected from 64 sites on September 1 and 13, 1993; from 63 sites on March 17 and 18, 1994; and from 64 sites on October 18 and 19, 1994. Discharge measurements help identify gains and losses of flow along stream channels (Riggs, 1972), and they aid in the comparison of low and high base-flow conditions. Specific-conductance and temperature measurements were used to help identify where ground-water discharged to the streams.

This report presents base-flow data collected during the course of the investigation. Gaining and losing reaches of the stream channels are delineated for low and high base-flow conditions. Sub-basins within the Little West Fork basin with surplus or deficient flow are also defined.

The Little West Fork basin is located in south-central Kentucky and north-central Tennessee. The study area consists of the surface-water drainage basin for the Little West Fork above site 50 (fig. 1). The Little West Fork basin above site 50 has a drainage area of 180 mi<sup>2</sup> which lies within parts of Trigg and Christian Counties, Kentucky, and parts of Stewart and Montgomery Counties, Tennessee (fig. 1). Most of the basin lies within the U.S. Army's Fort Campbell Military Reservation (fig. 1). The drainage areas for the most downstream sites along major tributaries to the Little West Fork include those for Noah Spring Branch (53.3 mi<sup>2</sup>), Dry Fork Creek (18.7 mi<sup>2</sup>), Piney Fork (50.2 mi<sup>2</sup>), and Fletchers Fork (24.3 mi<sup>2</sup>).

#### **BASE-FLOW DATA**

Base-flow data including discharge, temperature, and specific conductance were collected for the Little West Fork basin during low base-flow and high base-flow conditions. Data representing low base-flow conditions were collected on September 1 and 13, 1993, and October 18 and 19, 1994. Data representing high base-flow conditions were collected on March 17 and 18, 1994. Names and locations of base-flow measurement sites within the study area are shown in table 1. A hydrograph showing daily discharge measurements for the Little West Fork (site 40, fig. 1) and dates when base-flow data were collected is shown in figure 2.

#### **Discharge Data**

Discharge measurements were made at 64 of the 65 sites (60 stream sites and 4 springs) on September 1 and 13, 1993 (tables 2 and 3). The measurements were made during low base-flow conditions, and 45 of the measurement sites (70 percent) had zero flow. Most of the dry measurement sites lie in the head waters of Noah Spring Branch and Piney Fork Creek (fig. 1). Flows in the major streams in the basin were 4.38 ft<sup>3</sup>/s for Noah Spring Branch (site 16a), 4.02 ft<sup>3</sup>/s for Dry Fork Creek (site 16b), 0 ft<sup>3</sup>/s for Piney Fork (site 39), 0 ft<sup>3</sup>/s for Fletchers Fork (site 48), and 18.4 ft<sup>3</sup>/s for Little West Fork (site 50). Discharge measurements at sites with flow ranged from 0.02 ft<sup>3</sup>/s (site 22) to 18.4 ft<sup>3</sup>/s (site 50, the most downstream site in the basin). Measurement sites with discharge values for September 1 and 13, 1993, and dry, gaining, and losing reaches are shown in figure 3.

Discharge measurements were made at 63 of the 65 sites (59 stream sites and 4 springs) on March 17 and 18, 1994 (tables 4 and 5). One other spring was visited, but discharge could not be measured (table 5). Measurements were made during high base-flow conditions, and only 8 of the measurement sites (13 percent) had zero flow. Flows in the major streams in the basin were 124 ft<sup>3</sup>/s for Noah Spring Branch (site 16a), 66.3 ft<sup>3</sup>/s for Dry Fork Creek (site 16b), 58.5 ft<sup>3</sup>/s for Piney Fork (site 39), 28.7 ft<sup>3</sup>/s for Fletchers Fork (site 48), and 419 ft<sup>3</sup>/s for Little West Fork (site 50). Discharge measurements at sites with flow ranged from 0.01 ft $^3$ /s (sites 1, 14, and 29) to 419 ft $^3$ /s (site 50). Measurement sites with discharge values for March 17 and 18, 1994, and dry, gaining, and losing reaches are shown in figure 4.

Discharge measurements were made at 64 of the 65 sites (60 stream sites and 4 springs) on October 18 and 19, 1994 (tables 6 and 7). Measurements were again made during low base-flow conditions, and 44 of the measurement sites (69 percent) had zero flow. Most of the dry measurement sites lie in the head waters of Noah Spring Branch and Piney Fork Creek, as was observed in September 1993. Flows in the major streams in the basin were 4.69 ft<sup>3</sup>/s for Noah Spring Branch (site 16a), 4.98 ft<sup>3</sup>/s for Dry Fork Creek (site 16b), 0 ft<sup>3</sup>/s for Piney Fork (site 39), 0 ft<sup>3</sup>/s for Fletchers Fork (site 48), and 27.6 ft<sup>3</sup>/s for Little West Fork (site 50). Discharge measurements at sites with flow ranged from 0.03 ft<sup>3</sup>/s (sites 22 and 28) to 27.6 ft<sup>3</sup>/s (site 50). Measurement sites with discharge values for October 18 and 19, 1994, and dry, gaining, and losing reaches are shown in figure 5.

Discharge leaving the Little West Fork basin was greater during high base-flow conditions than during low base-flow conditions. Site 50 on the Little West Fork is the farthest downstream measurement site within the basin (fig. 1) and had the greatest discharge of all sites during all three measurement periods. Low base-flow measurements made at site 50 on September 13, 1993, and October 18, 1994, were 18.4 ft<sup>3</sup>/s (table 2) and 27.6 ft<sup>3</sup>/s (table 6), respectively. On March 17, 1994, during high base flow, the discharge measurement at site 50 was 419 ft<sup>3</sup>/s (table 4). Major tributaries on the south side of the study area (Piney Fork and Fletchers Fork) consistently had lower flow than the tributaries on the north side (Dry Fork Creek and Noah Spring Branch).

Discharge data were used to categorize stream reaches as gaining or losing, wet (little or no change in




Figure 1. Study area and locations of stream and spring measurement sites.

**Table 1**. Site and station numbers, station names, and locations of stream and spring measurement sites in the Little West Fork basin, Fort Campbell

	no. Station no. Station name		Location						
Site no.			Station no. Station name Latitude		de <sub>"</sub>	Longitud o '		ıde "	
1	03436370	Noah Spring Branch trib. near LaFayette, KY	36	38	39	87	36	51	
2	03436372	Noah Spring Branch trib. on Angels Rd. near Garrettsburg, KY	36	38	37	87	35	24	
3	03436373	Noah Spring Branch on Angels Rd. near Garretts- burg, KY	36	38	42	87	35	07	
4	03436375	Noah Spring Branch trib. off Palmyra Rd. near Garrettsburg, KY	36	38	19	87	33	26	
5	03436378	Unnamed trib. to Noah Cave at Lovelady Rd. near Garrettsburg, KY	36	39	46	87	34	28	
6	03436379	Unnamed trib. to Noah Cave on Boodie Rd. near Garrettsburg, KY	36	40	03	87	34	04	
7	03436382	Unnamed trib. to Noah Cave at Garrettsburg, KY	36	39	47	87	32	43	
8	03436384	Noah Cave near Garrettsburg, KY	36	38	47	87	33	21	
9	03436388	Noah Spring Branch at On the Line Rd. near Garrettsburg, KY	36	38	02	87	32	12	
10	03436390	Dry Fork Creek trib. near Army Airfield at Fort Campbell	36	40	57	87	29	03	
11	03436391	Dry Fork Creek trib. near Army Airfield at tower near Garrettsburg, KY	36	40	39	87	30	37	
12	03436394	Dry Fork Creek trib. at water tank near Army Airfield near Garrettsburg, KY	36	40	07	87	30	50	
13	03436396	Dry Fork Creek at Angels Rd. near Garrettsburg, KY	36	39	13	87	31	02	
14	03436397	Dry Fork Creek trib. at Angels Rd. near Garretts- burg, KY	36	39	19	87	30	22	
15	03436398	Dry Fork Creek trib. at Fort Campbell	36	38	06	87	28	33	
16	03436400	Noah Spring Branch	36	37	22	87	30	47	
16a			36	37	24	87	30	58	
16b			36	37	29	87	30	48	
17	03436401	Piney Fork at Destiny Trail near Legate, TN	36	34	51	87	42	24	

<sup>4</sup> Base-Flow Data for the Little West Fork Basin, Fort Campbell, Tennessee and Kentucky, 1993 and 1994

Table 1. Site and station numbers, station names, and locations of stream and spring measurement sites in the Little West Fork basin, Fort Campbell—Continued

					Lo	cation		
Site no.	Station no.	Station name	Latitude			Longitude o ' "		ude "
18	034364015	Piney Fork at Normandy Loop Rd. near Legate, TN	36	35	17	87	41	48
19	03436402	Piney Fork trib. at Normandy Loop Rd. near Legate, TN	36	35	17	87	41	48
20	03436403	Piney Fork Creek near Legate, TN	36	35	16	87	40	37
21	03436404	Piney Fork Creek at Rendevous Rd. near Legate, TN	36	35	16	87	39	54
22	03436405	Lake Kyle outflow at Destiny Trail near Legate, TN	36	34	58	87	39	24
23	03436406	Piney Fork Creek at Indian Mound Rd. near Legate, TN	36	35	22	87	38	46
24	03436407	Piney Fork trib. below Indian Mound Rd. near Legate, TN	36	35	24	87	38	44
25	03436408	Piney Fork above Jordan Spring Rd. near LaFayette, KY	36	35	38	87	37	49
26	03463409	Elk Fork Creek near Oakwood, TN	36	35	14	87	36	50
27	03437410	Elk Fork Creek at mouth near LaFayette, KY	36	35	38	87	37	47
28	034364104	Piney Fork near LaFayette, KY	36	36	22	87	37	56
29	034364105	Piney Fork trib. near LaFayette, KY	36	36	23	87	37	58
30	03436411	Piney Fork Creek north of Rose Hill Rd. near LaFayette, KY	36	37	06	<b>.</b> 87	36	38
31	03436412	Piney Fork at Engineer Rd. near Garrettsburg, KY	36	37	37	87	35	54
32	03436413	Piney Fork at Palmyra Rd. near Oakwood, TN	36	37	03	87	34	53
33	03436414	Piney Fork near Oakwood, TN	36	36	59	87	33	17
34	03436415	Piney Fork at mouth of Jordan Creek near Oakwood, TN	36	36	59	87	32	21
35	03436416	Jordan Creek at Jordan Spring Rd. near Oakwood, TN	36	35	40	87	33	16
36	03436417	Moss Creek at Palmyra Rd. near Oakwood, TN	36	35	53	87	34	37
37	03436418	Jordan Creek at Ghost Corps Trail near Oakwood, TN	36	36	29	87	32	42
38	03436419	Jordan Creek at mouth near Oakwood, TN	36	36	56	87	32	21


**Table 1.** Site and station numbers, station names, and locations of stream and spring measurement sites in the Little West Fork basin, Fort Campbell—Continued

	_				Lo	cation		
Site no.	Station no.	Station name		Station name Latitude				ıde "
39	03436420	Piney Fork	36	36	59	87	30	51
40	03436426	Little West Fork near Fort Campbell (East End Rd.)	36	36	38	87	28	11
41	03436430	Little West Fork Red River at Fort Campbell	36	36	28	87	26	57
42	03436433	Fletchers Fork near Woodlawn, TN	36	33	26	87	31	25
43	03436437	Fletchers Fork trib. near Woodlawn, TN	36	33	24	87	31	22
44	03436440	Fletchers Fork	36	34	27	87	30	42
45	03436441	Fletchers Fork on Woodlawn Rd. at walnut grove	36	35	19	87	27	57
46	03436442	Racoon Branch at walnut grove	36	35	17	87	27	56
47	03436443	Racoon Branch near walnut grove	36	34	31	87	27	57
48	03436444	Fletchers Fork near walnut grove	36	35	34	87	26	58
49	03436446	Fletchers Fork trib. on Even Rd. at Ringgold, TN	36	35	30	87	25	54
50	03436460	Little West Fork Red River near New Providence, TN	36	35	30	87	23	23
51	03436360	Noah Spring Branch on Angels Rd. near LaFayette, KY	36	38	24	87	39	15
52	03436361	Noah Spring Branch trib. on Angels Rd. near LaFayette, KY	36	38	21	87	39	01
53	03436362	Noah Spring Branch near LaFayette, KY	36	38	50	87	38	01
54	03436364	Noah Spring Branch trib. on HWY 107 near LaFayette, KY	36	40	42	87	38	30
55	03436365	Noah Spring Branch trib. at LaFayette Rd. near LaFayette, KY	36	39	02	87	38	00
56	03436366	Noah Spring Branch trib. near LaFayette, KY	36	38	50	87	38	00
57	03436368	Noah Spring Branch trib. on Angels Rd. near LaFayette, KY	36	38	19	87	37	18
58	03436369	Noah Spring Branch near LaFayette, KY	36	38	41	87	36	53
59	03436385	Noah Spring near Garrettsburg, KY	36	38	20	87	33	80

<sup>6</sup> Base-Flow Data for the Little West Fork Basin, Fort Campbell, Tennessee and Kentucky, 1993 and 1994

**Table 1.** Site and station numbers, station names, and locations of stream and spring measurement sites in the Little West Fork basin, Fort Campbell—Continued

								ation			
Site no.	Site no. Station no.	Station name	° L	atitu	de "	Lo	ngiti	nde.			
60	03436424	Dennis Spring to Little West Fork near Fort Campbell	36	36	36	87	29	44			
61	03436435	Unnamed spring on Old Dover Rd. at Woodlawn, TN	36	32	44	87	30	42			
62	03436445	Britton Spring at Ringgold, TN	36	35	35	87	25	56			
63		On the Line Rd. Spring to Noah Spring Branch	36	38	02	87	32	13			



Base-Flow Data for the Little West Fork Basin, Fort Campbell, Tennessee and Kentucky, 1993 and 1994

Table 2. Low base-flow data for streams in the Little West Fork basin, September 1 and 13, 1993

[ft³/s, cubic foot per second; °C, degrees Celsius;  $\mu S/cm$  at 25 °C, microSiemen per centimeter at 25 degrees Celsius]

Site no.	Station no.	Stream discharge, instantaneous (ft <sup>3</sup> /s)	Temperature (°C)	Specific conductance (µS/cm at 25 °C)
1	03436370	0		
2	03436372	0		
3	03436373	0		
4	03436375	0		
5	03436378	0		
6	03436379	0		
7	03436382	0		
8	03436384	0		
9	03436388	3.62	16.0	385
10	03436390	0		
11	03436391	0		
12	03436394	0		
13	03436396	3.50	17.1	375
14	03436397	0		
15	03436398	0		
<sup>a</sup> 16	03436400	8.40		
16a		4.38	18.2	382
16b		4.02	19.5	360
17	03436401	0		
18	034364015	0		
19	03436402	0		
20	03436403	0		<b></b>
21	03436404	0		
22	03436405	.02	25.3	184
23	03436406	0		
24	03436407	0		
25	03436408	0		
26	03463409	0		
27	03437410	0		
28	034364104	.08	19.8	245
29	034364105	0		
30	03436411	0		<del></del>
31	03436412	.13	21.2	128
32	03436413	.06	22.3	232
33	03436414	0		
34	03436415	0	<del></del>	
35	03436416	.39	20.3	371
36	03436417	0		
37	03436418	.13	19.2	347
38	03436419	0		
39	03436420	ő		
40	03436426	8.54	19.5	371
41	03436430	13.4	19.9	453
42	03436433	0		
43	03436437	0		

Table 2. Low base-flow data for streams in the Little West Fork basin, September 1 and 13, 1993-Continued

Site no.	Station no.	Stream discharge, instantaneous (ft <sup>3</sup> /s)	Temperature (°C)	Specific conductance (µS/cm at 25 °C)
44	03436440	0.64	19.1	331
45	03436441	0		
46	03436442	.08	17.6	407
47	03436443	0		
48	03436444	0		
49	03436446	0		
50	03436460	18.4	19.4	660
51	03436360	0		
52	03436361	0		
53	03436362	0		
54	03436364	0		
55	03436365	0		
56	03436366	0		
57	03436368	0		
58	03436369	0		

<sup>&</sup>lt;sup>a</sup> The streamflow measurement for this site was calculated by adding the measurements for sites 16a and 16b.

Table 3. Low base-flow data for springs in the Little West Fork basin, September 13, 1993 [ft<sup>3</sup>/s, cubic foot per second; °C, degrees Celsius; µS/cm at 25 °C, microSiemen per centimeter at 25 degrees Celsius]

Site no.	Station no.	Spring name	Discharge, instantaneous (ft <sup>3</sup> /s)	Temperature (°C)	Specific conductance (μS/cm at 25 °C)
59	03436385	Noah Spring	0.31	14.8	418
60	03436424	Dennis Spring	1.16	13.6	425
61	03436435	Unnamed spring	0		
62	03436445	Britton Spring	1.45	17.3	377




Figure 3. Little West Fork basin showing low base-flow stream and spring measurement sites and discharge measurements, September 1 and 13, 1993.

Table 4. High base-flow data for streams in the Little West Fork basin, March 17 and 18, 1994

[ft $^3$ /s, cubic foot per second; °C, degrees Celsius;  $\mu$ S/cm at 25 °C, microSiemen per centimeter at 25 degrees Celsius]

Site no.	Station no.	Stream discharge, instantaneous (ft <sup>3</sup> /s)	Temperature (°C)	Specific conductance (µS/cm at 25°C)
1	03436370	0.01	10.8	174
2	03436372	0		
3	03436373	13.1	11.2	222
4	03436375	56.1	10.4	228
5	03436378	1.29	12.7	216
6	03436379	1.06	9.9	125
7	03436382	.10	9.5	317
8	03436384	15.9	12.5	270
9	03436388	116	10.8	232
10	03436390	.06	10.0	67
11	03436391	.43	9.3	113
12	03436394	0		
13	03436396	45.2	12.8	280
14	03436397	.01	9.0	341
15	03436398	0		
<sup>a</sup> 16	03436400	190	10.0	252
16a		124		
16b	<u></u> .	· 66.3	<del>-</del> -	
17	03436401	.91	10.4	65
18	034364015	0		
19	03436402	2.09	9.6	60
20	03436403	5.80	9.3	54
21	03436404	5.28	8.4	55
22	03436405	1.85	11.3	57
23	03436406	8.97	8.4	57
24	03436407	.12	7.0	85
25	03436408	10.9	7.6	59
26	03463409	4.35	8.8	65
27	03437410	5.14	6.6	63
28	034364104	18.3	7.4	77
29	034364105	.01	7.4	191
30	03436411	22.4	10.2	105
31	03436412	24.7	10.8	110
32	03436413	36.8	10.5	113
33	03436414	38.4	9.7	121
34	03436415	5.49	10.1	88
35	03436416	10.4	14.0	164
36	03436417	1.82	9.0	45
37	03436418	15.4	9.0	146
38	03436419	9.20	9.6	150
39	03436420	58.5	8.0	138
40	03436426	284	10.0	248
41	03436430	317	10.0	242
42	03436433	0		
43	03436437	Ö		
44	03436440	18.8	13.0	140

<sup>12</sup> Base-Flow Data for the Little West Fork Basin, Fort Campbell, Tennessee and Kentucky, 1993 and 1994

**Table 4.** High base-flow data for streams in the Little West Fork basin, March 17 and 18, 1994—Continued

Site no.	Station no.	Stream discharge, instantaneous (ft <sup>3</sup> /s)	Temperature (°C)	Specific conductance (µS/cm at 25 °C)
45	03436441	31.9	8.0	165
46	03436442	6.09	**	180
47	03436443	2.14	7.0	75
48	03436444	28.7	12.0	170
49	03436446	0		
50	03436460	419	11.2	250
51	03436360	5.34	8.9	112
52	03436361	.42	9.8	249
53	03436362	10.5	10.2	160
54	03436364	1.29	6.6	
56	03436366	23.5	11.9	212
57	03436368	.32	8.4	66
58	03436369	38.2	11.5	202

<sup>&</sup>lt;sup>a</sup> The streamflow measurement for this site was calculated by adding the measurements for sites 16a and 16b.

Table 5. High base-flow data for springs in the Little West Fork basin, March 17, 1994

[ft³/s, cubic foot per second; °C, degrees Celsius; μS/cm at 25 °C, microSiemen per centimeter at 25 degrees Celsius]

Site no.	Station no.	Spring name	Discharge, instantaneous (ft <sup>3</sup> /s)	Temperature (°C)	Specific conductance (µS/cm at 25 °C)
59	03436385	Noah Spring	19.6	11.3	302
<sup>a</sup> 60	03436424	Dennis Spring			
61	03436435	Unnamed spring	0		
62	03436445	Britton Spring	12.2	12.0	270
63		On the Line Rd. Spring	4.83	12.6	211

<sup>&</sup>lt;sup>a</sup> A discharge measurement for this site was not possible during high base flow because the spring was submerged.




Figure 4. Little West Fork basin showing high base-flow stream and spring measurement sites and discharge measurements, March 17 and 18, 1994.

Table 6. Low base-flow data for streams in the Little West Fork basin, October 18 and 19, 1994

[ft $^3$ /s, cubic foot per second; °C, degrees Celsius;  $\mu$ S/cm at 25 °C, microSiemen per centimeter at 25 degrees Celsius]

Site no.	Station no.	Stream discharge, instantaneous (ft <sup>3</sup> /s)	Temperature (°C)	Specific conductance (µS/cm at 25 °C)
1	03436370	0		
2	03436372	0		
3	03436373	0		
4	03436375	0		
5	03436378	0		
6	03436379	0		
7	03436382	0		
8	03436384	0		
9	03436388	2.10	13.9	415
10	03436390	0		
11	03436391	0		
12	03436394	0		
13	03436396	3.91	14.4	380
14	03436397	0		
15	03436398	0		
<sup>a</sup> 16	03436400	9.67	15.2	309
16a		4.69		
16b		4.98		
17	03436401	0		
18	034364015	0		
19	03436402	0		
20	03436403	0		
21	03436404	0		**
22	03436405	.03	17.1	176
23	03436406	0		
24	03436407	0		
25	03436408	0		
26	03463409	0		
27	03437410	0		**
28	034364104	.03	16.8	276
29	034364105	0		
30	03436411	0		
31	03436412	.21	15.0	261
32	03436413	.20	16.8	299
33	03436414	.16	15.9	333
34	03436415	0		
35	03436416	.45	18.2	384
36	03436417	0		
37	03436418	.31	15.0	357
38	03436419	0		
39	03436420	0	***	
40	03436426	10.5	15.9	376
41	03436430	17.2	16.8	440
42	03436433	0		
43	03436437	Ö		
44	03436440	.80	15.2	337

Table 6. Low base-flow data for streams in the Little West Fork basin, October 18 and 19, 1994-Continued

Site no.	Station no.	Stream discharge, instantaneous (ft <sup>3</sup> /s)	Temperature (°C)	Specific conductance (μS/cm at 25 °C)
45	03436441	0		<del>-</del> -
46	03436442	.40	15.4	389
47	03436443	0		
48	03436444	0		
49	03436446	0		
50	03436460	27.6	16.3	393
51	03436360	0		
52	03436361	0		
53	03436362	0		
54	03436364	0		
55	03436365	0		
56	03436366	0		
57	03436368	0		
58	03436369	0		

<sup>&</sup>lt;sup>a</sup> The streamflow measurement for this site was calculated by adding the measurements for sites 16a and 16b.

Table 7. Low base-flow data for springs in the Little West Fork basin, October 18, 1994

[ft<sup>3</sup>/s, cubic foot per second; °C, degrees Celsius; µS/cm at 25 °C, microSiemen per centimeter at 25 degrees Celsius]

Site no.	Station no.	Spring name	Discharge, instantaneous (ft <sup>3</sup> /s)	Temperature (°C)	Specific conductance (µS/cm at 25 °C)
59	03436385	Noah Spring	0.04	14.1	413
60	03436424	Dennis Spring	.96	14.3	411
61	03436435	Unnamed spring	0		
62	03436445	Britton Spring	3.41	16.0	364




Figure 5. Little West Fork basin showing low base-flow stream and spring measurement sites and discharge measurements, October 18 and 19, 1994.

flow), dry, or unobserved. Gaining, losing, and dry stream reaches are bounded by one or more upstream and one downstream measurement sites. For this report, a gaining reach is defined as a reach in which flow has increased by at least 10 percent of its downstream discharge value. A losing reach is defined as a reach in which flow has decreased by at least 10 percent of its upstream discharge value. A wet reach is a reach in which the change in flow is less than 10 percent of its greater discharge value. A reach is categorized as dry if its upstream and downstream discharge measurements are 0 ft<sup>3</sup>/s. A reach not bounded by both upstream and downstream measurement sites is categorized as unobserved. More gaining stream reaches occurred during the high base-flow period (fig. 4) than either low base-flow period (fig. 3 and 5). More dry stream reaches occurred during the two low base-flow periods (fig. 3 and 5) than during the high base-flow period (fig. 4).

#### **Temperature Data**

Water temperature data also were collected at most of the sites that had flow when discharge measurements were taken. On September 1 and 13, 1993, water temperature data collected at 18 of the 19 sites with flow ranged from 13.6 to 25.3 °C (tables 2 and 3). On March 17 and 18, 1994, water temperature at 52 of the 56 sites with flow ranged from 6.6 to 14.0 °C (tables 4 and 5). On October 18 and 19, 1994, water temperature at 18 of the 20 sites with flow ranged from 13.9 to 18.2 °C (tables 6 and 7).

#### **Specific-Conductance Data**

Specific conductance also was measured at most of the sites that had flow. On September 1 and 13, 1993, specific conductance at 18 of the 19 sites with flow ranged from 128 to 660  $\mu$ S/cm (tables 2 and 3). On March 17 and 18, 1994, specific conductance at 52 of the 56 sites with flow ranged from 45 to 341  $\mu$ S/cm (tables 4 and 5). On October 18 and 19, 1994, specific conductance at 18 of the 20 sites with flow ranged from 176 to 440  $\mu$ S/cm (tables 6 and 7).

## AREAS OF SURPLUS OR DEFICIENT FLOW

Areas of surplus or deficient flow can be determined by comparing the change in flow of each subbasin with the flow of the entire basin. Such a comparison will reveal how much flow each sub-basin is contributing to the total flow of the basin. Areas of surplus or deficient flow were determined for low and high base-flow periods in the following manner:

- The drainage area for each sub-basin was measured from topographic maps.
- The change in flow for each sub-basin was calculated by subtracting the flow entering each sub-basin from the flow leaving each sub-basin. The change in flow was divided by the sub-basin drainage area. The change in flow per square mile for each sub-basin is shown in tables 8, 9, and 10.
- The flow per square mile of surface drainage for the entire study area was calculated by dividing the discharge at the most downstream site (50) (tables 2, 4, and 6) by the basin area at site 50 (180 mi<sup>2</sup>). Values of flow per square mile were calculated using discharge data collected on September 1 and 13, 1993, March 17 and 18, 1994, and October 18 and 19, 1994.
- The change in flow per square mile for each subbasin was compared to the flow per square mile for site 50. If the change in flow per square mile for a sub-basin was greater than 2 times or less than 1/2 of the flow per square mile for site 50, then it was defined as an area of surplus or deficient flow, respectively. Otherwise, the sub-basin was considered neither surplus nor deficient. Areas of surplus or deficient flow are shown in figures 6, 7, and 8.

Many areas of deficient flow occurred near the headwaters of the Little West Fork basin under all baseflow conditions. Fewer areas of deficient flow occurred near the mouth of the basin.

The flow per square mile for each major tributary basin in the study area was also calculated (table 11). The values of flow per square mile for the Noah Spring Branch and Dry Fork Creek basins in the northern part of the study area were greater than those for the Piney Fork and Fletchers Fork basins in the southern part of the study area under all base-flow conditions. The tributary basin areas in the southern part of the study area had deficient flow (0 ft<sup>3</sup>/s/mi<sup>2</sup>) for both low base-flow periods and neither surplus nor deficient flow for the high base-flow period. All

**Table 8.** Low base-flow data for drainage areas in the Little West Fork basin, September 1 and 13, 1993

[mi<sup>2</sup>, square mile; ft<sup>3</sup>/s, cubic foot per second; ft<sup>3</sup>/s/mi<sup>2</sup>, cubic foot per second per square mile; change in flow per square mile, positive numbers reflect gain and negative numbers reflect loss]

Site and sub-basin no. (figure 6)	Sub-basin area (mi <sup>2</sup> )	Discharge entering sub- basin (ft <sup>3</sup> /s)	Discharge leaving sub- basin (ft <sup>3</sup> /s)	Change in flow per square mile (ft <sup>3</sup> /s/mi <sup>2</sup> )
1	0.15	0	0	0
2	.46	0	0	0
3	2.39	0	0	0
4	1.76	0	0	0
5	4.77	0	0	0
6	1.99	0	0	0
7	2.36	0	0	0
8	5.17	0	0	0
9	1.64	0	3.62	2.21
10	.11	0	0	0
11	3.88	0	0	0
12 13	.23	0	0	0
15 14	5.45	0	3.50	.64
15	1.11 . <b>95</b>	0 0	0 0	0 0
16a	.93 1.00	3.62	4.38	.76
16b	6.99	3.50	4.02	.76 .07
17	2.60	0	0	.07 0
18	2.88	0	0	0
19	.26	ő	0	0
20	5.81	ő	Ö	0
21	.93	ŏ	Ö	ő
22	2.07	Õ	.02	.01
23	1.63	.02	0	01
24	.47	0	0	0
25	2.23	0	0	0
26	7.01	0	0	0
27	.97	0	0	0
28	.90	0	.08	.09
29	.47	0	0	0
30	2.00	.08	0	04
31	.84	0	.13	.15
32	.96	.13	.06	07
33	4.50	.06	0	01
34	.72	0	0	0
35	4.50	0	.39	.09
36	3.12	0	0	0
37	2.33	.39	.13	11
38	1.21	.13	0	11
39	1.77	0	0	0
40	5.85	8.40	8.54	.02
41	2.65	8.54	13.4	1.83
42	9.73	0	0	0

Table 8. Low base-flow data for drainage areas in the Little West Fork basin, September 1 and 13, 1993—Continued

Site and sub-basin no. (figure 6)	Sub-basin area (mi <sup>2</sup> )	Discharge entering sub- basin (ft <sup>3</sup> /s)	Discharge leaving sub- basin (ft <sup>3</sup> /s)	Change in flow per square mile (ft <sup>3</sup> /s/mi <sup>2</sup> )
43	0.42	0	0	0
44	2.83	0	.64	.23
45	4.60	.64	0	14
46	1.27	0	.08	.06
47	3.94	0	0	0
48	1.50	.08	0	05
49	.88	0	0	0
50	23.7	13.4	18.4	.21
51	14.8	0	0	0
52	1.44	0	0	0
53	2.29	0	0	0
54	1.92	0	0	0
55	8.96	0	0	0
56	.10	0	0	0
57	1.03	0	0	0
58	1.16	0	0	0
				.10
Entire basin	180	0	18.4	(Flow per square mile)

**Table 9.** High base-flow data for drainage areas in the Little West Fork basin, March 17 and 18, 1994

[mi<sup>2</sup>, square mile; ft<sup>3</sup>/s, cubic foot per second; ft<sup>3</sup>/s/mi<sup>2</sup>, cubic foot per second per square mile; change in flow per square mile, positive numbers reflect gain and negative numbers reflect loss]

Site and sub-basin no. (figure 7)	Sub-basin area (mi <sup>2</sup> )	Discharge entering sub- basin (ft <sup>3</sup> /s)	Discharge leaving sub- basin (ft <sup>3</sup> /s)	Change in flow per square mile (ft <sup>3</sup> /s/mi <sup>2</sup> )
1	0.15	0	0.01	0.07
2	.46	0	0	0
3	2.39	38.2	13.1	-10.5
4	1.76	13.1	56.1	24.4
5	4.77	0	1.29	.27
6	1.99	0	1.06	.53
7	2.36	0	.10	.04
8	5.17	2.45	15.9	2.60
9	1.64	56.1	116	36.5
10	.11	0	.06	.54
11	3.88	.06	.43	.10
12	.23	0	0	0
13	5.45	.43	45.2	8.22
14	1.11	0	.01	.01
15	.95	0	0	0
16a	1.00	116	124	8.00
16b	6.99	45.2	66.3	3.02
17	2.60	43.2 0		
			.91	.35
18	2.88	.91	0	32
19	.26	0	2.09	8.04
20	5.81	2.09	5.80	.64
21	.93	5.80	5.28	56
22	2.07	0	1.85	.89
23	1.63	7.13	8.97	1.13
24	.47	0	.12	.26
25	2.23	9.09	10.9	.81
26	7.01	0	4.35	.62
27	.97	4.35	5.14	.81
28	.90	16.0	18.3	2.51
29	.47	0	.01	.02
30	2.00	18.3	22.4	2.04
31	.84	22.4	24.7	2.74
32	.96	24.7	36.8	12.6
33	4.50	36.8	38.4	.36
34	.72	38.4	5.49	-45.7
35	4.50	0	10.4	2.31
36	3.12	0	1.82	.58
37	2.33	12.2	15.4	1.36
38	1.21	0	9.20	-5.12
39	1.77	14.7	58.5	24.7
40	5.85	248	284	6.07
41	2.65	284	317	12.4
42	9.73	0	0	0
		6.6		/ 1

Table 9. High base-flow data for drainage areas in the Little West Fork basin, March 17 and 18, 1994—Continued

Site and sub-basin no. (figure 7)	Sub-basin area (mi <sup>2</sup> )	Discharge entering sub- basin (ft <sup>3</sup> /s)	Discharge leaving sub- basin (ft <sup>3</sup> /s)	Change in flow per square mile (ft <sup>3</sup> /s/mi <sup>2</sup> )
44	2.83	0	18.8	6.64
45	4.60	18.8	31.9	2.85
46	1.27	2.14	6.09	3.11
47	3.94	0	2.14	.54
48	1.50	38.0	28.7	-6.19
49	.88	0	0	0
50	23.7	346	419	3.09
51	14.8	0	5.34	.36
52	1.44	0	.42	.29
53	2.29	5.76	10.5	2.07
54	1.92	0	1.29	.67
56	9.06	1.29	23.5	2.45
57	1.03	0	.32	.31
58	1.16	34.3	38.2	3.34
				2.33
Entire basin	180	0	419	(Flow per
				square mile)

**Table 10.** Low base-flow data for drainage areas in the Little West Fork basin, October 18 and 19, 1994

[mi<sup>2</sup>, square mile; ft<sup>3</sup>/s, cubic foot per second; ft<sup>3</sup>/s/mi<sup>2</sup>, cubic foot per second per square mile; change in flow per square mile, positive numbers reflect gain and negative numbers reflect loss]

Site and sub-basin no. (figure 8)	Sub-basin area (mi <sup>2</sup> )	Discharge entering sub- basin (ft <sup>3</sup> /s)	Discharge leaving sub- basin (ft <sup>3</sup> /s)	Change in flow per square mile (ft <sup>3</sup> /s/mi <sup>2</sup> )
1	0.15	0	0	0
2	.46	0	0	0
3	2.39	0	0	0
4	1.76	0	0	0
5	4.77	0	0	0
6	1.99	0	0	0
7	2.36	0	0	0
8	5.17	0	0	0
9	1.64	0	2.10	1.28
10	.11	0	0	0
11	3.88	0	0	0
12	.23	0	0	0
13	5.45	0	3.91	.72
14	1.11	0	0	0
15	.95	0	0	0
16a	1.00	2.10	4.69	2.59
16b	6.99	3.91	4.98	.15
17	2.60	0	0	0
18	2.88	0	0	0
19	.26	0	0	0
20	5.81	0	0	0
21	.93	0	0	0
22	2.07	0	.03	.01
23	1.63	.03	0	02
24	.47	0	0	0
25	2.23	0	0	0
26	7.01	0	0	0
27	.97	0	0	0
28	.90	0	.03	.03
29	.47	0	0	0
30	2.00	.03	0	02
31	.84	0	.21	.25
32	.96	.21	.20	01
33	4.50	.20	.16	01
34	.72	.16	0	22
35	4.50	0	.45	.10
36	3.12	0	0	0
37	2.33	.45	.31	06
38	1.21	.31	0	26
39	1.77	0	0	0
40	5.85	9.67	10.5	.14
41	2.65	10.5	17.2	2.53
42	9.73	0	0	0
				~

**Table 10.** Low base-flow data for drainage areas in the Little West Fork basin, October 18 and 19, 1994—Continued

Site and sub-basin no. (figure 8)	Sub-basin area (mi <sup>2</sup> )	Discharge entering sub- basin (ft <sup>3</sup> /s)	Discharge leaving sub- basin (ft <sup>3</sup> /s)	Change in flow per square mile (ft <sup>3</sup> /s/mi <sup>2</sup> )
44	2.83	0	0.80	0.28
45	4.60	.80	0	17
46	1.27	0	.40	.32
47	3.94	0	0	0
48	1.50	.40	0	27
49	.88	0	0	0
50	23.7	17.2	27.6	.44
51	14.8	0	0	0
52	1.44	0	0	0
53	2.29	0	0	0
54	1.92	0	0	0
55	8.96	0	0	0
56	.10	0	0	0
57	1.03	0	0	0
58	1.16	0	0	0
Entire basin	180	0	27.6	.15 (Flow per square mile)

Table 11. Flow per square mile for major tributary basins in the Little West Fork basin

[mi<sup>2</sup>, square mile; ft<sup>3</sup>/s/mi<sup>2</sup>, cubic foot per second per square mile]

	Site	Basin area	Flow per square mile (ft <sup>3</sup> /s/mi <sup>2</sup> )		
Basin	no.	(mi <sup>2</sup> )	September 1993	March 1994	October 1994
Tributary					
Noah Spring Branch	16a	53.3	0.08	2.33	0.09
Dry Fork Creek	16b	18.7	.21	3.55	.27
Piney Fork	39	50.2	0	1.17	0
Fletchers Fork	48	24.3	0	1.18	0
Entire basin Little West Fork	50	180	.10	2.33	.15




Figure 6. Little West Fork basin showing areas of surplus or deficient flow during low base flow, September 1 and 13, 1993.




Figure 7. Little West Fork basin showing areas of surplus or deficient flow during high base flow, March 17 and 18, 1994.




Figure 8. Little West Fork basin showing areas of surplus or deficient flow during low base flow, October 18 and 19, 1994.

tributary basin areas in the northern part of the study area had neither surplus nor deficient flow for all base-flow periods, with the exception of the Dry Fork Creek basin, which had surplus flow of 0.21 ft<sup>3</sup>/s/mi<sup>2</sup> for September 1993 (table 11).

#### **SUMMARY**

The Little West Fork drainage basin is located in south-central Kentucky and north-central Tennessee. Most of the basin lies within the Fort Campbell Military Reservation. Base-flow data including discharge, temperature, and specific conductance were collected from 65 sites within the basin. Low base-flow data were collected from 64 sites on September 1 and 13, 1993, and from 64 sites on October 18 and 19, 1994. On September 1 and 13, 1993, discharge at sites with flow ranged from 0.02 to 18.4 ft<sup>3</sup>/s at the most downstream site in the basin. On October 18 and 19, 1994, discharge at sites with flow ranged from 0.03 to 27.6 ft<sup>3</sup>/s at the most downstream site in the basin. High base-flow data were collected from 63 sites on March 17 and 18, 1994. Discharge at sites with flow ranged from 0.01 to 419 ft<sup>3</sup>/s at the most downstream site in the basin. Discharge from the basin was greater during high base-flow conditions than during low base-flow conditions. Major tributaries on the south side of the study area (Piney Fork and Fletchers Fork) consistently had lower flow than the tributaries on the north side (Dry Fork Creek and Noah Spring Branch).

Discharge data were used to categorize stream reaches and sub-basins. Stream reaches were characterized as gaining or losing, wet, dry, or unobserved for each base-flow measurement period. More gaining stream reaches occurred during the high base-flow period than either low base-flow period. More dry stream reaches occurred during the two low base-flow periods than during the high base-flow period.

Sub-basin areas with surplus or deficient flow were defined within the basin. Drainage areas for each stream measurement site were delineated and measured from topographic maps. Change in flow per square mile for each sub-basin was calculated using data from each base-flow measurement period. The calculated values were used to define the areas of surplus or deficient flow for low and high base-flow conditions. Many areas of deficient flow occurred near the headwaters of the Little West Fork basin under all base-flow conditions. Fewer areas of deficient flow occurred near the mouth of the basin.

The flow per square mile for each major tributary basin in the study area was also calculated. The values of flow per square mile for the Noah Spring Branch and Dry Fork Creek basins in the northern part of the study area were greater than those for the Piney Fork and Fletchers Fork basins in the southern part of the study area under all base-flow conditions. The tributary basin areas in the southern part of the study area had deficient flow (0 ft³/s/mi²) for both low base-flow periods and neither surplus nor deficient flow for the high base-flow period. All tributary basin areas in the northern part of the study area had neither surplus nor deficient flow for all base-flow periods, with the exception of the Dry Fork Creek basin, which had surplus flow for September 1993.

#### REFERENCES CITED

Carey, M.S., 1990, The delineation of karst drainage to identify possible contaminant migration routes from the Campbell Army Airfield, Fort Campbell, Kentucky (M.S. thesis): Richmond, Kentucky, Eastern Kentucky University, 61 p.

Riggs, H.C., 1972, Low-flow investigations: Techniques of Water-Resources Investigations of the U.S. Geological Survey, Book 4, Chap. B1, p. 11.